

FACTS ABOUT FENTANYL TOXICITY AND HOW WE CAN HELP REDUCE THE MORTALITY

Zoha Jabbar *

Department of Forensic Science, Faculty of Medicine and Allied Health Sciences, The Islamia University of Bahawalpur, Punjab, Pakistan

Submitted: July 20, 2024; Revised: December 14, 2024; Accepted: December 22, 2024; Published: April 30, 2025

ABSTRACT

Due to its part in the opioid epidemic, fentanyl, a powerful synthetic opioid drug, has received a lot of attention recently. It is estimated that this medicine, which was initially created to treat pain, has 50–100 times the potency of morphine. Especially in cancer patients and following surgery, fentanyl's potent effects make it beneficial in alleviating severe pain. Approximately 50–100 times more strong than morphine and 50 times more potent than heroin, fentanyl is a synthetic opioid analgesic. It was initially created for medical usage as a potent painkiller, often taken via injection, transdermal patches, or lozenges for severe pain control, such as after surgery or in patients with cancer. Fentanyl is an effective treatment for severe pain in a medical environment. To patients with a high tolerance for opioids, it is given by medical specialists in amounts that are carefully monitored. Overdose and other harmful effects are kept to a minimum thanks to the regulated environment. Specific medical situations, such as surgery or intensive care, are ideal for its quick onset and brief duration of effect. Fentanyl's illicit use is a problem for the drug. Overdose deaths from illicitly produced fentanyl have increased globally, frequently being combined with other narcotics like heroin or cocaine. Due to the drug's strength and erratic distribution in street narcotics, users could unintentionally take deadly dosages. An alarming increase in opioid-related fatalities has resulted from this, creating a severe public health disaster. Even though fentanyl is essential in medicine for treating pain, it can be lethal when used illicitly. A concerning rise in overdose fatalities has been caused by its extraordinary strength and extensive use in the illegal drug trade. Stricter regulation, harm reduction techniques, increased access to addiction treatment, and mental health services are just a few of the many initiatives needed to address the fentanyl problem.

Keywords: Fentanyl, Toxicity, Antidot, Opioids.

*Corresponding Author. E-mail: zohajabbar46@gmail.com

INTRODUCTION

Dr. Paul Janssen, a Belgian pharmacologist and the creator of Janssen Pharmaceutical, created fentanyl for the first time in 1960. Fentanyl was created by the research team when they were looking for novel analgesic (pain-relieving) substances. The medication was first marketed under the trade name "Sublimaze," and it quickly became well-liked in medical settings due to its potent pain-relieving abilities. Due to its quick onset and brief duration of action, it was predominantly utilized as an intravenous anesthetic during surgical operations. This made it appropriate for usage during surgeries and other medical treatments. Following its discovery, pharmaceutical firms created a number of fentanyl formulations to enhance its medical uses. These included sublingual pills, nasal sprays, transdermal patches, and lozenges (lollipops), all of which provided a regulated and long-lasting delivery

of the painkiller [1]. Fentanyl has been used medically for the treatment of severe pain, particularly when other opioids have failed to relieve the pain or have had unpleasant side effects. Fentanyl did, however, have hazards due to its tremendous potency, particularly when misused. An upsurge in overdoses and fatalities from opioid-related causes has been brought on by the drug's abuse and unauthorized distribution. Fentanyl is still an essential drug for the treatment of pain, even with the problems brought on by overuse, especially in medical settings where its strength and quick-acting characteristics may be successfully managed and monitored.

STRUCTURE AND CLASSIFICATION

The group of medications known as opioids includes fentanyl. It is a synthetic opioid, which means it is produced chemically as opposed to being sourced



from a natural source like opium. Fentanyl is a derivative of phenylpiperidine in terms of structure. The compound is known chemically as N-(1-phenethylpiperidin-4-yl)-N-phenylpropanamide. Fentanyl's molecular weight is about 336.47 g/mol, and its chemical formula is $C_{22}H_{28}N_2O$ [2].

As an opioid analgesic, fentanyl is extremely strong due to its structural features. Its potency allows for efficient pain relief at low dosages, but it also raises the possibility of overdose and other negative effects, especially when taken incorrectly or without a doctor's supervision. Due to its extreme dependency and significant potential for misuse, fentanyl and its analogs are listed as Schedule II restricted narcotics in the United States. Their medicinal usage is closely regulated, and only licensed healthcare practitioners may prescribe them for valid therapeutic objectives, such as controlling extreme pain in carefully monitored medical environments.

SYNTHESIS

It is discussed how the parent opioid fentanyl and its analogs may be synthesized in alternative and more efficient ways. After conducting optimization experiments for each synthetic step, the pathways described demonstrate high-yielding transformations that lead to these potent analgesics. The basic three-step approach gave rise to a panel of four fentanyls with outstanding yields (73–78%) coupled with their more prevalent hydrochloride and citric acid salts [3].

PHARMACOKINETICS (PK)

The study of a drug's absorption, distribution, metabolism, and excretion (ADME) is referred to as pharmacokinetics (PK). For a medicine like fentanyl to be effective and safe when used medically, it is essential to understand its PK profile.

Absorption

Transdermal patches, nasal sprays, lozenges, and intravenous injections are just a few of the different ways that fentanyl is administered. Fentanyl has a speedy beginning of effect because to its rapid and full absorption when given intravenously. Transdermal patches deliver the medication via the skin in a regulated manner, maintaining stable blood levels for a long time. Different formulations also show different rates of absorption.

Distribution

High lipophilicity of fentanyl enables fast passage of the blood-brain barrier and distribution throughout the central nervous system (CNS). It is also delivered to many bodily tissues. Because of its high affinity for opioid receptors, it has strong analgesic effects.

Metabolism

Fentanyl's major metabolic route takes place in the liver, where the enzyme cytochrome P450 3A4 (CYP3A4) converts it into inactive metabolites, mostly norfentanyl [4]. It's also possible for certain small metabolites to arise.

Elimination

The urine is the main elimination route for fentanyl and its metabolites. Depending on the formulation and method of administration, fentanyl's elimination half-life might change. For instance, although transdermal patches may have a half-life of roughly 17 hours, intravenous fentanyl generally has a half-life of around 2 to 4 hours. It is essential to remember that due to fentanyl's strength and quick beginning of effect, medical personnel must administer and monitor the drug with caution. Due to its tremendous strength, there is a chance of overdose and severe negative consequences, especially when taken outside of proper medical settings or in conjunction with other drugs. When using fentanyl or any other medicine, always abide by the authorized dosage and administration guidelines provided by a licensed healthcare professional [5].

MOA(MECHANISM OF ACTION)

Fentanyl's main mechanism of action (MOA) is how it interacts with opioid receptors in the central nervous system (CNS). The major action of fentanyl as an opioid analgesic is the binding of certain opioid receptors, namely the mu-opioid receptors.

Simplified explanation of MOA

Fentanyl works as an agonist, which means it binds to and activates mu-opioid receptors that are spread throughout the brain and spinal cord. The perception and processing of pain signals are facilitated by these receptors, which are a component of the body's natural pain modulation system.

Inhibition of pain signaling

Once fentanyl attaches to mu-opioid receptors, it prevents the nervous system from sending pain signals. By reducing the feeling of pain, this activity has strong analgesic (pain-relieving) benefits.

Modulation of reward pathways

Fentanyl has painkilling properties in addition to modulating reward pathways in the brain, which results in sensations of pleasure and relaxation. This feature of its MOA increases the likelihood that it will be abused and become addictive.

Respiratory depression

Fentanyl and other opioids have the principal adverse effect of respiratory depression. Breathing can be slowed down by the activation of opioid receptors in the brainstem, which can lower the sensitivity of the respiratory centers. When taken with other central nervous system depressants or at greater doses, this impact can be harmful. It's critical to understand that fentanyl requires cautious prescribing and monitoring by medical experts due to its powerful analgesic qualities, as well as the risk of abuse and overdose. Because of the drug's tremendous strength and potential for side effects, it is often only used to treat really severe pain [6].

NORMAL/TOXIC DOSE /POISONING

The amount of fentanyl that is "normal" or safe to take varies greatly depending on the formulation, the

patient's health, and their tolerance to opioids. Given how powerful fentanyl is as an opioid, even little variations in dosage can have negative effects. As a result, it must only be used as directed by a licensed healthcare provider. Due to its strength, fentanyl is normally supplied in micrograms (mcg) for legitimate medical usage. For instance, intravenous dosages for surgical anesthetic or acute pain control may vary from 25 to 100 mcg. Fentanyl can be released from transdermal patches over a period of 72 hours at rates of 12 to 100 mcg/hour. The limited therapeutic index of fentanyl, however, makes it critical to recognize when talking about hazardous dosages or poisoning. Toxic respiratory depression, coma, or death can result from a relatively little dosage increase. Fentanyl used for medicinal purposes improperly or illegally can be quite hazardous [7]. According to variables including tolerance, body weight, and unique opioid sensitivity, the fatal dose of fentanyl might differ significantly from person to person. Fentanyl can be fatal to people who are not opioid-tolerant even at doses as little as 2 milligrams. It is critical to understand that non-pharmaceutical fentanyl, which is frequently discovered in illicit substances or fake pharmaceuticals, can be far more strong and unpredictable, increasing the danger of overdose and poisoning [8]. Call emergency services right away and seek immediate medical assistance if you believe someone has been exposed to fentanyl and is exhibiting overdose signs including difficulty breathing, excessive sleepiness, or unconsciousness. Again, to guarantee correct dosage and lower the danger of side effects or poisoning, only take fentanyl under the supervision and prescription of a licensed healthcare practitioner.

PLASMA LEVEL DETERMINATION

Especially when it is used for pain relief or anesthesia, determining the plasma level of fentanyl is a crucial component of its therapeutic usage. Monitoring fentanyl levels in the blood enables medical personnel to give patients an appropriate dosage while reducing the danger of side effects, including respiratory depression. Blood samples from the patient are taken at predetermined intervals after the medicine has been delivered in order to determine the plasma level of fentanyl. Depending on the delivery method and the particular formulation being utilized, the time of the blood draws will vary. Blood samples are often obtained quickly after a medication injection for intravenous delivery in order to evaluate the drug's fast start of action and peak plasma concentration [10]. Blood samples may be obtained at various intervals to track the steady-state plasma levels for transdermal patches or other formulations with a protracted release. The findings of the plasma level determination are utilized for therapeutic drug monitoring (TDM) and, if necessary, to change the dose of the fentanyl. Making ensuring the

medication stays within the therapeutic range, where it can effectively treat pain without having any negative side effects, is the aim. An extremely strong opioid, fentanyl should be used under the direct supervision of trained medical specialists. When performing pain management or anesthetic operations, one of the instruments utilized to optimize fentanyl administration and guarantee patient safety is plasma level determination [11].

SIGNS AND SYMPTOMS

A strong opioid, fentanyl can cause a variety of indications and symptoms when taken improperly or abusively as well as when it is used as prescribed under a doctor's care. Based on the user's tolerance, the dosage, and the method of administration, fentanyl's effects might vary. The following are some typical fentanyl usage symptoms and warning signs: Pain alleviation is one of fentanyl's main effects because it is an opioid analgesic. In medical contexts, it is frequently administered to treat extremely painful conditions. Fentanyl, like other opioids, has the ability to elicit sensations of euphoria, calm, and relaxation. The possibility for drug misuse is increased by these side effects. Fentanyl has the potential to make users feel sleepy and drowsy, which will impair their ability to focus and stay attentive. Breathing becomes shallow and slowly as a result of respiratory depression, one of the most serious adverse effects of fentanyl and other opioids. Lack of oxygen might result in respiratory arrest in extreme circumstances. Fentanyl usage can result in gastrointestinal problems that can make people feel sick to their stomachs. Opioids, such as fentanyl, can constrict the pupils. Fentanyl has the potential to fog the mind, impair judgment, and impair cognitive function. Fentanyl and other opioids can cause sluggish bowel motions, which can result in constipation. Fentanyl's adverse effects might cause itching or a rash in some people. Fentanyl lowers blood pressure, causing lightheadedness and dizziness. Taking too much fentanyl may result in an overdose, which is potentially fatal. Insanity, severe respiratory depression, and blue lips or fingertips are indications of an overdose [12]. It is crucial to understand that fentanyl's strength and risk of overdose make it highly dangerous to abuse or consume unsupervised. A considerable proportion of opioid-related overdose deaths have been linked to illicitly acquired fentanyl, particularly when it is combined with other narcotics or sold as counterfeit medication.

CLINICAL USES

Fentanyl is a strong opioid drug with several therapeutic applications, especially for pain control and anesthesia. It is a useful tool in medical settings due to its high potency and rapid beginning of effect. Fentanyl is often used in the multiple therapeutic settings. Fentanyl is frequently used to treat severe pain, particularly when other opioids or analgesic

drugs are ineffective. It can be given intravenously, topically, nasally, topically, transdermally, or intrathecally. During surgical operations, fentanyl is routinely used as a supplement to general Anesthetic. During and after surgery, it may assist to deliver analgesia and lessen discomfort. Fentanyl may be used in long-acting formulations, such as transdermal patches, [13] for those with chronic pain disorders, such as cancer-related pain or other incapacitating illnesses, to provide long-lasting pain relief. Fentanyl may be administered during birth in some circumstances to relieve pain for women who are unable to have an epidural anesthetic. Fentanyl is used for procedural sedation and analgesia during several medical procedures that call for sedation and pain relief, such as endoscopy or cardiac catheterization. Patients who have had surgery may occasionally take fentanyl to treat their pain in the postoperative phase [14]. Fentanyl can be used to treat pain and sedate patients who are mechanically ventilated in intensive care units (ICUs) and other critical care settings. In emergency situations, such as those involving trauma or recent injuries, fentanyl can be used to quickly relieve severe pain. The fact that fentanyl is a strong opioid with a significant risk of respiratory depression and other adverse effects must be noted. Its usage ought to be properly regulated and managed by licensed medical personnel. To reduce the danger of overdose or unfavorable responses, fentanyl dosage and delivery need to be closely watched. Fentanyl also has a significant potential for misuse and dependency, underscoring the significance of administering it safely and with the proper medical supervision.

DEATH RATE IN DIFFERENT YEARS

The opioid was present in 75% of the over 92,000 drug overdose deaths in 2020. A drug overdose has claimed the lives of more than 932,000 individuals since 1999.1 In the United States, 91,799 people died from drug overdoses in 2020. From 2019 (21.6 per 100,000) to 2020 (28.3 per 100,000), the age-adjusted rate of overdose fatalities increased by 31%.The leading cause of drug overdose deaths at the moment is opioids, particularly synthetic opioids (other than methadone). Synthetic opioids were present in 82.3% of overdose fatalities involving opioids. In 2020, 68,630 drug overdose fatalities (74.8% of all drug overdose deaths) involved opioids. Both with and without the use of synthetic opioids, drug overdose fatalities involving psychostimulants like methamphetamine are rising

DEATH RATE IN DIFFERENT COUNTRIES

Over 10% of the world's countries have drug overdose mortality rates that are higher than four per 100,000 people, including Russia and Libya (where rates are down) as well as Libya (where rates are rising). Data from five developed countries with significant drug mortality rates were compared, and it was found that rates in the United States and Canada had lately accelerated and alarmingly

increased (this is likely due to the introduction of fentanyl and rising stimulant usage). The categorization of some suicides as accidents may be to blame for a purported recent drop in Canada's drug suicide rate. Verifying accuracy and improving knowledge of drug overdose deaths and ways to avoid them are key priorities.

PRODUCTS AVAILABILITY

Fentanyl is now accessible in a number of pharmaceutical formulations, mostly for medical use under stringent control and prescription by licensed healthcare professionals as of my most recent knowledge update in September 2021 [15]. Among the popular fentanyl products are: Fentanyl is available in intravenous (IV) formulations for use during surgery and other medical procedures that call for strong and quick pain management. Transdermal Patches: Fentanyl patches provide a regulated delivery of the drug via the skin over a number of days, making them useful for long-term pain treatment. Fentanyl nasal sprays are used to provide quick pain relief, especially when IV administration is impractical. Patients who are unable to take drugs orally occasionally utilize fentanyl lozenges or lollipops for pain management [16]. Fentanyl sublingual pills are intended to dissolve under the tongue and offer prompt pain relief. It's crucial to keep in mind that local laws and nation differences affect where fentanyl products can be found. Fentanyl is a prohibited chemical in many jurisdictions because of its extreme potency and potential for misuse. Its usage in medicine is therefore strictly regulated. Fentanyl that is produced and supplied illegally is a serious public health issue since it increases the danger of overdose and death and is frequently discovered in illegal substances or fake prescriptions. Illegal and extremely dangerous to public safety, the availability of illicit fentanyl goods.

ANTIDOTS

Naloxone, often known by the brand name Narcan, is the substance that serves as an antidote for fentanyl and other opioids. Because naloxone blocks the actions of opioids like fentanyl on opioid receptors in the brain and central nervous system, it can swiftly reverse the effects of these drugs. In cases of opioid overdose, particularly fentanyl overdose, naloxone can be given right away and quickly bring about a return to normal breathing. In emergency cases where an opioid overdose is suspected, it is a life-saving drug and a crucial tool. Naloxone is simple to use even for those who are not medical experts because it is offered in a variety of formulations, including injectables and nasal sprays [17]. Numerous first responders, members of law enforcement, and healthcare professionals have received training in how to treat opioid overdose victims using naloxone. If you have access to naloxone and are trained to use it, administer it if you believe someone is suffering from an opioid

overdose, including one from fentanyl, and emergency medical assistance is on the way. But always keep in mind that naloxone is not a replacement for skilled medical assistance, and anybody who has taken naloxone should still seek emergency medical attention for additional assessment and treatment.

HOW TO CONTROL ITS CONSUMPTION

Controlling fentanyl use is essential for combating the opioid crisis and advancing public health and safety. The following tactics can aid in reducing fentanyl consumption: Before prescribing fentanyl and other opioids, medical providers should thoroughly assess the necessity for them. Establishing precise regulations for the prescription of opioids, including fentanyl, will help guarantee that these drugs are only used when necessary and in the right amounts [18]. Monitoring tools like prescription medication monitoring programs can aid in finding instances of possible overprescribing or abuse. Individualized instruction about the dangers and advantages of fentanyl should be given to patients who have been prescribed it. There should be detailed instructions on how to use the product, any possible adverse effects, and the significance of sticking to the recommended dosages. Promoting the use of non-opioid pain reduction techniques including physical therapy, non-opioid drugs, and alternative therapies can help lessen the need for powerful opioids like fentanyl. Implementing harm reduction initiatives can help avoid opioid overdoses and lessen the negative effects of fentanyl usage. These initiatives include needle exchange programs and availability to naloxone. To stop the drug from being diverted to the black market and other unlawful sources, it is crucial to strictly regulate and enforce the laws surrounding fentanyl's manufacturing, distribution,

and illegal use. Public awareness campaigns can assist in educating the general public about the risks associated with the abuse of fentanyl and the possibility of an overdose. Information on how to spot and handle opioid overdoses may also be found in these advertisements [19]. Access to addiction treatment and support services should be improved so that people who are dependent on opioids, especially fentanyl, may get the aid they need to rehabilitate and cut back on use. Ongoing studies of opioid analgesics, painkiller substitutes, and addiction therapies may help us better understand opioid-related problems and develop approaches that are more successful. To successfully manage the fentanyl pandemic and solve the larger opioid crisis, it is imperative to develop a holistic strategy that requires cooperation between healthcare professionals, decision-makers, law enforcement, community groups, and the general public [20].

CONCLUSION

In conclusion, even while fentanyl is essential for treating medical pain, its potential for abuse and the alarming increase in overdose fatalities necessitate a comprehensive response. This entails tighter controls on its manufacturing and distribution, enhanced harm reduction and addiction treatment initiatives, as well as greater public education and awareness campaigns. In the world of medicine and drug policy, it is still difficult to strike a balance between the advantages of fentanyl and the need to lessen the hazards that go along with it.

Author(s) Contribution: resources and editing, J. Zoha.

Funding: No funding involved.

Ethical Approval: Not applicable

Conflict of Interest: Nil.

Consent for Publication: All authors approved the manuscript for publication.

REFERENCES

1. Jaffe J. H., Martin W. R. Opioid analgesics and antagonists. Pharmacological Basis of Therapeutics 7th Ed., A. G. Gilman, L. S. Goodman, T. W. Rail, F. Murad. MacMillan, New York 1985; 517.
2. Physicians Desk Reference 48th ed. Medical Economics Data Production Company, Montvale 1994; 1092–1093.
3. Marshall B. E., Wollman H. General anesthetics. Pharmacological Basis of Therapeutics 7th ed., A. G. Gilman, L. S. Goodman, T. W. Rail, F. Murad. MacMillan, New York 1985; 276–301.
4. McClain D. A., Hug C. C., Jr. Intravenous fentanyl kinetics. Clin Pharmacol Therap 1980; 28: 106–114.
5. Hess P., Steibler G., Herz A. Pharmacokinetics of fentanyl in man and rabbit. Eur J Clin Pharmacol 1972; 4: 137–141.
6. Mather L. E. Clinical pharmacokinetics of fentanyl and its newer derivatives. Clin Pharmacokinet 1983; 8: 422–446.
7. Bower S., Hull C. J. Comparative pharmacokinetics of fentanyl and alfentanil. Br J Anaesth 1982; 54: 871–877.
8. Bower S. The uptake of fentanyl by erythrocytes. J Pharm Pharmacol 1982; 34: 181–185.
9. Fung D. L., Eisele J. H. Fentanyl pharmacokinetics in awake volunteers. J Clin Pharmacol 1980; 20: 652–658.
10. Bovill J. G., Sebel P. S. Pharmacokinetics of high-dose fentanyl. A study in patients undergoing cardiac surgery. Br J Anaesth 1980; 52: 795–801.
11. Stoeckel H., Hengstmann J. H., Schuttler J. Pharmacokinetics of fentanyl as a possible explanation for recurrence of respiratory depression. Br J Anaesth 1979; 51: 741–745.
12. Koren G., Goresky G., Crean P., Klein J., MacLeod S. M. Pediatric fentanyl dosing based on pharmacokinetics during cardiac surgery. Anesth Analg 1984; 63: 577–582.
13. Cartwright P., Prys-Roberts C., Gill K., Dye A., Stafford M., Gray A. Ventilatory depression related to plasma fentanyl concentrations during and after anesthesia in humans. Anesth Analg 1983; 62: 966–974.
14. Henderson G. L. The fentanyl. Service Training and Education. American Association for Clinical Chemistry. 1990; Vol 12: 7–14.
15. Kumar K., Morgan D. J. Determination of fentanyl and alfentanil in plasma by high-performance liquid chromatography with ultraviolet detection. J Chromatogr 1987; 419: 464–468.
16. Calis K. A., Kohler D. R., Corso D. M. Transdermal administered fentanyl for pain management. Clin Pharm 1992; 11: 22–36.
17. Jones D. B., et al. Two unusual deaths associated with the fentanyl transdermal drug delivery system. 45th Ann. Meeting American Academy Forensic Sciences Program. Boston, MA 1993; 169.
18. Marquardt K. A., Tharratt R. S. Inhalation abuse of fentanyl patch. J Toxicol Clin Toxicol 1994; 32: 75–78.
19. Streisand J. B., Varvel J. R., Stanski D. R., et al. Absorption and bioavailability of oral transmucosal fentanyl citrate. Anesthesiology 1991; 75: 223–229.
20. Stanley T. H., Hague B., Mock D. L., et al. Oral transmucosal fentanyl citrate (lollipop) premedication in human volunteers. Anesth Analg 1989; 69: 21–27.