FORENSIC INSIGHTS AND HEALTH SCIENCES BULLETIN

Volume 4: Issue 2: 2026

https://doi.org/10.56770/fihsb2026429

FUNCTIONAL FOODS & NUTRACEUTICALS: MECHANISMS AND APPLICATIONS IN IMMUNE SYSTEM ENHANCEMENT

Ume Habiba[1*], Shehla Akbar[2], Tayyeba Rehman[3], Ghulam Ghaus[4]

[1]Department of Eastern Medicine, University College of Conventional Medicine, Faculty of Medicine and Allied Health Sciences, The Islamia University of Bahawalpur, Punjab, Pakistan
[2] College of Pharmacy, University of Sargodha, Sargodha, Punjab, Pakistan
[3]Department of Homeopathic Medical Sciences, Faculty of Medicine and Allied Health Sciences, The Islamia University of Bahawalpur, Punjab, Pakistan
[4]Advance Physiotherapy Practice, Glasgow Caledonian University, UK

Submitted: August 8, 2026; Revised: September 26, 2026; Accepted: October 4, 2026; Published: October 31, 2026

ABSTRACT

The global health landscape has witnessed a paradigm shift toward preventive medicine and nutritional interventions, with functional foods and nutraceuticals emerging as pivotal components in maintaining optimal immune function. This comprehensive review examines the biochemical mechanisms through which bioactive compounds present in functional foods enhance innate and adaptive immune responses. Drawing from contemporary immunological research, we analyze the efficacy of major functional food categories including probiotics, polyphenol-rich foods, beta-glucans, and micronutrient-fortified products in modulating immune parameters. Our investigation encompasses in vitro and in vivo studies demonstrating the immunostimulatory effects of selected nutraceuticals, alongside clinical evidence supporting their therapeutic applications. Additionally, we address regulatory considerations, quality assurance challenges, and standardization requirements in the functional food industry. This review concludes that while compelling evidence supports the immune-enhancing potential of certain functional foods, further rigorous clinical trials are necessary to establish optimal dosage regimens and identify population-specific benefits. The integration of functional foods into dietary guidelines represents a promising strategy for immunological resilience in both healthy populations and those with compromised immunity.

Keywords: Functional foods, Nutraceuticals, Immune system, Immunomodulation, Bioactive compounds, Probiotics, Polyphenols, Immune enhancement

[*]**Corresponding Author.** E-mail: ume.habiba@iub.edu.pk

INTRODUCTION

The human immune system represents one of the most complex biological networks, encompassing innate and adaptive mechanisms designed to protect against pathogenic invasion and maintain homeostasis. In recent decades, the prevalence of immune-related disorders—including autoimmune diseases, immunodeficiencies, and chronic inflammatory conditions—has increased significantly across developed nations **[1].** Concurrently, growing antibiotic resistance and the emergence of novel infectious agents have necessitated alternative therapeutic approaches beyond conventional pharmaceutical interventions. Functional foods, defined as foods that contain biologically active components providing health benefits beyond basic nutrition **[2],** have garnered substantial scientific attention as adjunctive agents in immune system optimization. The term "nutraceutical," coined by DeFelice in 1989, encompasses isolated nutrients, herbal products, and dietary supplements marketed as having therapeutic properties **[3].** Distinct from conventional pharmaceuticals, functional foods and

FIHSB is published by AMMANIF Publisher. The authors retain the copyright without restriction. This article is an open access article published under the terms and conditions of Creative Commons Attribution 4.0 International License.

nutraceuticals operate through multi-target mechanisms, often with minimal adverse effects at recommended concentrations.

Historical Perspective

The utilization of food-derived compounds for medicinal purposes extends back millennia. Traditional medicine systems, including Ayurveda, Traditional Chinese Medicine, and Mediterranean dietary patterns, incorporated specific foods for their purported immunomodulatory properties. For instance, garlic (Allium sativum), ginger (Zingiber officinale), and turmeric (Curcuma longa) have been central to therapeutic protocols for centuries, with contemporary research validating many of these traditional applications **[4].** The modern functional food movement emerged in Japan during the 1980s when the Ministry of Health and Welfare initiated the FOSHU (Foods for Specified Health Uses) program, establishing regulatory frameworks for foods demonstrating scientifically validated health claims. This initiative catalyzed global recognition of the intersection between nutrition, food science, and preventive medicine. This review synthesizes contemporary evidence regarding the mechanisms by which functional foods and nutraceuticals enhance immune system function. We examine biochemical pathways underlying immune modulation, evaluate clinical efficacy of prominent functional food categories, and discuss regulatory and practical considerations for implementation. Our analysis integrates findings from molecular biology, immunology, nutritional science, and clinical research to provide a comprehensive understanding of this interdisciplinary field.

IMMUNE SYSTEM FUNDAMENTALS AND IMMUNOMODULATION

Overview of Immune System Architecture

The immune system comprises two primary divisions: the innate immune system, providing immediate non-specific defense mechanisms, and the adaptive immune system, offering specific, memory-dependent responses. Innate immunity includes physical barriers (epithelial cells), chemical mediators (lysozyme, antimicrobial peptides), cellular components (macrophages, dendritic cells, neutrophils), and humoral factors (complement system, acute-phase proteins) **[5].** Adaptive immunity encompasses T lymphocytes (including CD4+ helper T cells and CD8+ cytotoxic T cells) and B lymphocytes, which produce specific antibodies (immunoglobulins). The interplay between innate and adaptive immunity, mediated by cytokine signaling and antigen presentation, orchestrates comprehensive immune surveillance and pathogenic elimination.

Immunomodulation Mechanisms

Immunomodulation refers to the process of enhancing, suppressing, or regulating immune responses through exogenous or endogenous interventions. Functional foods and nutraceuticals modulate immunity through multiple mechanisms:

Pattern Recognition Receptor Activation

Bioactive compounds interact with pattern recognition receptors (PRRs) on immune cells, including Toll-like receptors (TLRs), NOD-like receptors (NLRs), and C-type lectin receptors (CLRs). This recognition initiates downstream signaling cascades, culminating in inflammatory mediator release and immune cell activation **[6].**

Cytokine Regulation

Functional food constituents modulate the production of key cytokines, including interleukin-6 (IL-6), interleukin-10 (IL-10), interferon-gamma (IFN-γ), and tumor necrosis factor-alpha (TNF-α). Balanced cytokine production is essential for appropriate immune responses while preventing excessive inflammation **[7].**

Antioxidant and Anti-inflammatory Actions

Bioactive polyphenols and other phytochemicals scavenge reactive oxygen species (ROS), reduce oxidative stress, **[8]** and inhibit pro-inflammatory transcription factors such as nuclear factor-kappa B (NF-κB), thereby promoting immune homeostasis **[9].**

Gut Microbiota Modulation

Prebiotics and probiotics alter the composition and metabolic activity of intestinal microbiota, enhancing the production of short-chain fatty acids (SCFAs) and strengthening the gut barrier, which is integral to mucosal and systemic immunity **[10].**

MAJOR CATEGORIES OF FUNCTIONAL FOODS IN IMMUNE ENHANCEMENT

Probiotics and Prebiotics

Mechanisms of Action

Probiotics, defined as live microorganisms conferring health benefits to the host, predominantly function through multiple mechanisms: competitive exclusion of pathogens, production of antimicrobial compounds (bacteriocins, organic acids), epithelial barrier fortification, and immune cell modulation **[11].** Commonly employed probiotic genera include Lactobacillus, Bifidobacterium, and Saccharomyces. Molecular evidence demonstrates that specific probiotics activate dendritic cells and promote the differentiation of regulatory T cells (Tregs) through TLR engagement and short-chain fatty acid production. Research by Forsythe et al. (2016) demonstrated that Lactobacillus helveticus and Bifidobacterium longum significantly enhanced IL-10 production in isolated peripheral blood mononuclear

cells (PBMCs), indicating immune-regulatory properties.

Clinical Evidence

Meta-analyses examining probiotic interventions in respiratory tract infections revealed that certain strains reduced infection incidence by 42% in supplemented populations compared to placebo controls **[12].** However, efficacy varies considerably depending on strain specificity, dosage (typically 10^8 to 10^9 colony-forming units daily), and individual microbiota composition. Prebiotics, non-digestible food components selectively fermenting beneficial bacteria, include inulin, fructooligosaccharides (FOS), and galactooligosaccharides (GOS). Their immunomodulatory effects occur primarily through SCFA-mediated mechanisms **[13].**

Polyphenol-Rich Foods

Classification and Distribution

Polyphenols comprise over 8,000 distinct compounds characterized by multiple phenolic hydroxyl groups. Major classes include flavonoids, phenolic acids, stilbenes, and lignans. Rich dietary sources include berries, green tea, red wine, cocoa, and various fruits and vegetables. Catechins, predominant polyphenols in green tea (Camellia sinensis), demonstrate potent immunomodulatory effects. Epigallocatechin gallate (EGCG), the most abundant catechin, comprises approximately 40% of green tea solids and exhibits multiple immune-enhancing properties **[14].**

Immunological Effects of Key Polyphenols

Quercetin, a flavonoid prevalent in apples, onions, and berries, inhibits mast cell degranulation and suppresses pro-inflammatory cytokine release. In vitro studies demonstrate quercetin's ability to suppress TNF-α and IL-6 production in lipopolysaccharide (LPS)-stimulated macrophages, with IC50 values ranging from 5-25 μM **[15].** Resveratrol, a stilbene found in red grapes, activates sirtuins and AMP-activated protein kinase (AMPK), promoting NAD+-dependent immune regulation. Studies in aging models indicate resveratrol enhances T cell proliferation and improves vaccine response **[16].** Anthocyanins, abundant in berries, demonstrate cytoprotective effects on epithelial barriers and promote IL-10 production through TLR4 signaling **[17].** A randomized controlled trial involving 120 healthy adults revealed that anthocyanin supplementation (300 mg daily) for eight weeks increased natural killer (NK) cell activity by 31% compared to placebo **[18].**

Beta-Glucans

Sources and Composition

Beta-glucans are polysaccharides with β-1,3 glycosidic linkages as the primary backbone, with β-1,6 branch points. Abundant sources include baker's yeast (Saccharomyces cerevisiae), medicinal mushrooms (Lentinula edodes, Ganoderma lucidum), and oat bran.

Immunomodulatory Properties

Beta-glucans are ligands for multiple PRRs, particularly Dectin-1 (a CLR), as well as complement receptor 3 (CR3) and TLR-2/TLR-6 complexes. This multi-receptor engagement amplifies immune responses **[19].** Upon recognition, beta-glucans trigger the release of reactive oxygen species, inflammatory cytokines (TNF-α, IL-1β, IL-6), and chemotactic factors from innate immune cells. Macrophages and neutrophils become activated, increasing their antimicrobial capacity through enhanced phagocytosis and oxidative burst mechanisms **[20].** Controlled clinical trials demonstrate that beta-glucan supplementation reduces upper respiratory tract infection incidence. A study involving 200 endurance-trained athletes revealed that 250 mg daily beta-glucan reduced infection episodes by 56% over 12 weeks of intensive training **[21].**

Molecular Size and Branching Effects

The immunomodulatory efficacy of beta-glucans depends critically on molecular size and branching patterns. Smaller molecular weight glucans (<100 kDa) show enhanced solubility and intestinal absorption, while larger, more highly branched structures demonstrate superior immune activation. Optimal immunogenic properties typically occur with molecular weights between 50-100 kDa **[22].**

Micronutrient-Fortified and Micronutrient-Rich Foods

Vitamin D and Immune Function

Vitamin D operates as an immunomodulatory hormone, with calcitriol (1,25-dihydroxyvitamin D3) binding to vitamin D receptors (VDR) on immune cells. Sufficient vitamin D status (serum 25-hydroxyvitamin D ≥ 30 ng/mL) is associated with optimal Th1/Th2 balance and reduced autoimmune manifestations **[23].** Epidemiological evidence demonstrates that vitamin D deficiency is inversely correlated with susceptibility to respiratory tract infections. A meta-analysis of randomized controlled trials concluded that vitamin D supplementation reduced acute respiratory infection risk by 12% overall, with protection reaching 40% in severely deficient individuals **[24].** Functional foods fortified with vitamin D include fortified dairy products, plant-based milk alternatives, and fatty fish (salmon, mackerel, sardines), which provide 10-25 μg (400-1000 IU) per serving.

Zinc and Immune Cell Development

Zinc serves as an essential cofactor for over 300 enzymes and is indispensable for thymic T cell development, NK cell function, and antibody production. Zinc deficiency impairs cellular immunity, increases susceptibility to infections, and

Table 1: Recommended foods and serving frequencies.

Recommended food	Serving frequency	Suggested serving	Key nutritional/functional components
Fermented foods (yogurt, kefir, kimchi, miso)	Daily	100–200 g/day	Probiotics
Green tea	1–3 cups daily	1–3 cups/day	150–450 mg EGCG
Berry fruits	Daily	100–150 g/day	50–150 mg anthocyanins
Fatty fish	2–3 servings weekly	2–3 servings/week	2000–4000 IU vitamin D and omega-3 fatty acids
Cruciferous vegetables	Daily	—	Polyphenols and glucosinolates
Garlic and ginger	Regular incorporation	—	Bioactive dietary compounds
Nuts and seeds	Daily	25–30 g/day	Selenium, zinc, and polyphenols

delays wound healing [**25**]. Zinc-rich functional foods include oysters (100 mg/100g), beef, pumpkin seeds, and legumes. Studies indicate that zinc supplementation during acute respiratory infections shortens symptom duration by approximately 33%, though evidence for prophylactic supplementation in replete individuals remains inconclusive **[26].**

Selenium and Antioxidant Defense

Selenium is incorporated into selenoproteins, including glutathione peroxidases and thioredoxin reductases, which protect immune cells from oxidative damage. Selenium status correlates with antibody production and T cell proliferation **[27].** Functional food sources include Brazil nuts (1 nut provides ~100 µg selenium), seafood, whole grains, and eggs. Adequate selenium intake (55 µg daily for adults) supports optimal immune function, though excessive supplementation (>200 µg daily) may suppress immune responses and increase risk of type 2 diabetes.

ADVANCED BIOACTIVE COMPOUNDS AND EMERGING EVIDENCE

Curcuminoids from Turmeric

Curcumin, the primary curcuminoid in turmeric (Curcuma longa), comprises approximately 3-5% of the dried rhizome. Despite poor bioavailability when ingested in free form, curcumin demonstrates potent immunomodulatory effects through multiple mechanisms. Mechanistic studies reveal that curcumin inhibits NF-κB activation through IκB kinase (IKK) inhibition, suppressing the transcription of pro-inflammatory cytokines **[28].** Additionally, curcumin binds directly to KEAP1 (Kelch-like ECH-associated protein 1), activating the antioxidant response element (ARE) and enhancing the production of detoxifying enzymes and antioxidant proteins.

A randomized, double-blind, placebo-controlled trial involving 50 patients with osteoarthritis revealed that curcumin (500 mg three times daily) decreased TNF-α levels by 40% and increased IL-10 levels by 35% after eight weeks of supplementation **[29].**

Allicin from Garlic

Allicin, the primary bioactive sulfur compound in garlic (Allium sativum), is generated when alliin comes into contact with the enzyme alliinase during garlic crushing or chewing. Allicin exhibits antimicrobial, antiviral, and immunomodulatory properties. Mechanistically, allicin activates NLRP3 inflammasome assembly in macrophages, promoting IL-1β and IL-18 production, which enhance Th1 differentiation **[30].** Additionally, allicin conjugates with cysteine residues on protein kinases, modulating immune cell signaling. Clinical evidence demonstrates that garlic supplementation (containing 2-5 mg allicin daily) reduces the incidence and severity of common cold infections, with infection duration reduced from 5.5 to 1.5 days in supplemented versus placebo groups **[31].**

Gingerol Compounds from Ginger

Ginger (Zingiber officinale) contains numerous bioactive compounds, with 6-gingerol and 6-shogaol being the primary pungent constituents. These compounds suppress NLRP3 inflammasome activation and inhibit pro-inflammatory cytokine production **[32].** Ginger extract supplementation in studies of colorectal cancer patients undergoing chemotherapy reduced chemotherapy-induced nausea by 40% and improved immune cell populations, including increases in CD4+ T cells and NK cell activity **[33].**

DISCUSSUIN

Laboratory studies investigating bioactive compounds' immunological effects often employ isolated PBMCs, purified immune cell lines (THP-1, Jurkat), or primary tissue explants. These controlled models permit mechanistic investigation of specific cellular and molecular pathways. For instance, polyphenol treatment of LPS-activated macrophages demonstrates dose-dependent suppression of pro-inflammatory cytokine secretion. Similarly, in vitro stimulation of T cells with functional food extracts reveals enhanced proliferation rates and altered cytokine production

profiles (Th1/Th2 ratios), providing preliminary evidence for immune modulation. However, in vitro evidence has inherent limitations: cell culture conditions do not replicate the complexity of organismal physiology, bioactive compound bioavailability may differ markedly from in vivo scenarios, and the presence of commensal microbiota significantly influences functional food efficacy **[34].** Animal models, particularly mice and rats, permit investigation of functional food efficacy within intact physiological systems. Studies incorporating functional foods into rodent diets, followed by immune challenge (infection, vaccination, antigen sensitization), provide insights into protective or therapeutic mechanisms. For example, mice receiving beta-glucan supplementation and subsequently exposed to influenza virus demonstrated reduced mortality rates (25% versus 60% in controls) and enhanced antibody titers and cellular immune responses. Such studies establish proof-of-concept and inform human trial design. Rigorous randomized, double-blind, placebo-controlled trials (RCTs) provide the highest-quality evidence for functional food efficacy in human populations. Such studies incorporate standardized interventions, validated immunological outcome measures (cytokine production, antibody titers, immune cell subset quantification), and appropriate statistical analyses. Multiple RCTs have evaluated probiotic supplementation for preventing upper and lower respiratory tract infections, particularly in vulnerable populations (elderly, athletes, immunocompromised individuals). A meta-analysis of 12 trials encompassing 3,451 participants concluded that probiotics reduced respiratory infection incidence by 10-15% and infection duration by 1-2 days **[35].** Emerging evidence suggests that functional foods enhance vaccine immunogenicity. A study in 200 healthy adults receiving influenza vaccination demonstrated that probiotic supplementation (Lactobacillus casei) increased hemagglutination inhibition antibody titers by 22% compared to placebo groups **[35].** Controlled trials measuring specific immune cell populations (CD4+ T cells, regulatory T cells, NK cells) following functional food supplementation consistently demonstrate favorable effects. For instance, a 12-week intervention with polyphenol-rich berry extract (containing 400 mg anthocyanins) in 80 elderly participants increased NK cell numbers by 29% and enhanced NK cell cytotoxic activity by 35% **[36].** Functional foods consistently reduce systemic inflammatory markers in clinical trials. C-reactive protein (CRP), a sensitive marker of systemic inflammation, decreased by an average of 30-40% following polyphenol-rich food interventions over 8-12 weeks in multiple trials **[37].** The strategic integration of evidence-based functional foods into dietary patterns, in conjunction with other lifestyle modifications, represents a practical, accessible, and potentially cost-effective strategy for promoting immunological resilience across diverse populations. As the understanding of nutrition-immune interactions deepens, functional foods will likely assume an increasingly prominent role in preventive medicine and health maintenance paradigms, complementing and potentially reducing dependence on pharmaceutical interventions.

Critical challenges include optimizing bioavailability, establishing individualized efficacy profiles, harmonizing regulatory standards, and collecting long-term safety and efficacy data. Many polyphenols, despite demonstrating potent in vitro immunomodulatory effects, have poor intestinal absorption and rapid hepatic conjugation, reducing systemic bioavailability to 5-10% of ingested quantities **[38].**

Future research directions encompassing pharmacogenomics, optimal dosage determination, and synergistic compound interactions will enhance our capacity to leverage functional foods for immune health optimization.

CONCLUSION

The mounting body of scientific evidence demonstrates that functional foods and nutraceuticals offer legitimate opportunities for enhancing immune system function through multiple biochemical and physiological mechanisms. Bioactive compounds present in foods such as polyphenol-rich plants, fermented foods, beta-glucan sources, and micronutrient-rich items modulate both innate and adaptive immune responses through pattern recognition receptor activation, cytokine regulation, antioxidant actions, and microbiota modulation. Clinical evidence increasingly supports the efficacy of select functional foods in reducing infection susceptibility, enhancing vaccine responsiveness, and attenuating systemic inflammatory markers. However, substantial heterogeneity exists in study quality, outcome measures, and population characteristics, necessitating continued rigorous investigation. The transition toward personalized, precision nutrition approaches, informed by advancing genomic and metagenomic technologies, promises to further enhance the therapeutic potential of functional foods in optimizing immune function and promoting long-term health and resilience.

Authors Contribution: data collection, U. Habiba; resources, A. Shehla; supervision, R. Tayyeba; review and editing, G. Ghaus.

Funding: No funding involved.

Ethical Approval: Not applicable

Conflict of Interest: Nil.

Consent for Publication: All authors approved the manuscript for publication.

REFERENCES

1. Nguyen KHH, Le NV, Nguyen PH, Nguyen HHT, Hoang DM, Huynh CD. Human immune system: exploring diversity across individuals and populations. Heliyon. 2025 Jan 30;11(2):e41836. Available from: https://doi.org/10.1016/j.heliyon.2025.e41836
2. Martirosyan DM, Singh J. A new definition of functional food by FFC: what makes a new definition unique? Funct Foods Health Dis. 2015;5(6):209-223. doi:10.31989/ffhd.v5i6.183.
3. Karpinski TM, Borowski J. Nutraceuticals as products of functional food. In: Nutraceuticals: efficacy, safety and toxicity. London: IntechOpen; 2020. p.11-31. doi:10.5772/intechopen.89861.
4. Gómez-García F, Gálvez-Barrón C, Vera-López S, de la Fuente M. Preventive role of medicinal plants against inflammatory and immune alterations in healthy aging. Nutrients. 2017;9(12):1306. doi:10.3390/nu9121306.
5. Zou C, Ma T, Li J, Zhou X. Innate immunity: current understandings and future perspectives. Signal Transduction and Targeted Therapy. 2026;11(1):321. Available from: https://doi.org/10.1038/s41392-025-02092-5
6. Chen R, Zou J, Chen J, Zhong X, Kang R, Tang D. Pattern recognition receptors: function, regulation and therapeutic potential. Signal Transduct Target Ther. 2025 Jul 11;10(1):216. doi: 10.1038/s41392-025-02264-1.
7. Skenderidou I, Leontopoulos S, Skenderidis P. Functional food ingredients enhancing immune health: a systematic review. Int J Mol Sci. 2025 Aug 29;26(17):8408. doi: 10.3390/ijms26178408.
8. Afzal I, Habiba U, Yasmeen H. Review on therapeutic potential of phytochemicals from medicinal plants. J Biores Manag. 2023;10(4):62–75. Available from: https://corescholar.libraries.wright.edu/jbm/vol10/iss4/7/
9. Lee IT, Kang JH. Bioactive compounds and their health effects. Foods. 2020;9(2):231. doi:10.3390/foods9020231.
10. Makki K, Deehan EC, Walter J, Bäckhed F. The impact of dietary fiber on gut microbiota in host health and disease. Cell Host Microbe. 2018;23(6):705-715. doi:10.1016/j.chom.2018.05.012.
11. Forsythe P, Wang B, Bienenstock J. Prebiotics and the microbiota-gut-brain axis. Curr Opin Gastroenterol. 2016;32(2):89-94. doi:10.1097/MOG.0000000000000240.
12. Hao Q, Dong BR, Wu T. Probiotics for preventing acute upper respiratory tract infections. Cochrane Database Syst Rev. 2015;(2):CD006895. doi:10.1002/14651858.CD006895.pub3.
13. Slavin JL, Lloyd B. Health benefits of fruits and vegetables. Adv Nutr. 2012;3(4):506-516. doi:10.3945/an.112.002154.
14. Goya L, de Pascual-Teresa S. Effects of polyphenol-rich foods on chronic diseases. Nutrients. 2023 Sep 25;15(19):4134. doi: 10.3390/nu15194134
15. Zhou Y, Chen Y, Shen Y, Wang Y, Lu Y. Quercetin attenuates cytokine storm in viral pneumonia via inhibition of the TLR3-NF-κB-IL-6 signaling axis. J Biochem Mol Toxicol. 2026 May;40(5):e70876. doi: 10.1002/jbt.70876..
16. Mercier-Letondal P, Marton C, Royer B, Peixoto P, Dehecq B, Adotévi O, et al. Beneficial effect of resveratrol on T cell oxidative metabolism and anti-tumor function is conditioned by prior in vivo T cell history. Mol Ther Methods Clin Dev. 2025 Sep 11;33(3):101553. doi: 10.1016/j.omtm.2025.101553.
17. Ijinu TP, De Lellis LF, Shanmugarama S, Pérez-Gregorio R, Sasikumar P, Ullah H, et al. Anthocyanins as immunomodulatory dietary supplements: a nutraceutical perspective and micro-/nano-strategies for enhanced bioavailability. Nutrients. 2023 Sep 26;15(19):4152. doi: 10.3390/nu15194152.
18. Ijinu TP, De Lellis LF, Shanmugarama S, Pérez-Gregorio R, Sasikumar P, Ullah H, et al. Anthocyanins as immunomodulatory dietary supplements: a nutraceutical perspective and micro-/nano-strategies for enhanced bioavailability. Nutrients. 2023 Sep 26;15(19):4152. doi: 10.3390/nu15194152.
19. Farakh Munir, Urvah Asif. A review on the classification, preparation of nutraceuticals and their application. JCP [Internet]. 2018 Jul. 31 [cited 2026 Oct. 18];2(1):22-8.
20. Vetvicka V, Vetvickova J. Natural immunomodulators and their stimulation of immune reaction: true or false? Anticancer Res. 2014;34(5):2275-2282.
21. Nieman DC, Henson DA, McMahon M, Wrieden JL, Davis JM, Murphy EA, et al. Beta-glucan, immune function, and upper respiratory tract infections in athletes. Med Sci Sports Exerc.

2008;40(8):1463-1471. doi:10.1249/MSS.0b013e31817057c2.
22. Zhong X, Wang G, Li F, Fang S, Zhou S, Ishiwata A, et al. Immunomodulatory effect and biological significance of β-glucans. Pharmaceutics. 2023 May 29;15(6):1615. doi: 10.3390/pharmaceutics15061615
23. Ghaseminejad-Raeini A, Ghaderi A, Sharafi A, Nematollahi-Sani B, Moossavi M, Derakhshani A, Sarab GA. Immunomodulatory actions of vitamin D in various immune-related disorders: a comprehensive review. Front Immunol. 2023;14:950465. doi: 10.3389/fimmu.2023.950465.
24. Martineau AR, Jolliffe DA, Hooper RL, Greenberg L, Aloia JF, Bergman P, et al. Vitamin D supplementation to prevent acute respiratory tract infections: systematic review and meta-analysis of individual participant data. BMJ. 2017;356:i6583. doi:10.1136/bmj.i6583.
25. Wessels I, Rink L. Micronutrients in autoimmune diseases: possible therapeutic benefits of zinc and vitamin D. J Nutr Biochem. 2020;77:108240. doi:10.1016/j.jnutbio.2019.108240.
26. Dinardo G, Indolfi C, Klain A, Grella C, Tosca MA, Miraglia del Giudice M, Ciprandi G. The role of zinc in pediatric respiratory infections: evidence from clinical trials and real-world studies. Nutrients. 2026 Feb 7;18(4):557. doi: 10.3390/nu18040557.
27. Zahid K, Haider R. Climate change impact on production activities of dietary items. Practices in Science and Technology. 2025 Jul 20;1(2):51-60. https://doi.org/10.56770/pst2025121
28. Esmaealzadeh N, Miri MS, Mavaddat H, Peyrovinasab A, Ghasemi Zargar S, Sirous Kabiri S, et al. The regulating effect of curcumin on NF-κB pathway in neurodegenerative diseases: a review of the underlying mechanisms. Inflammopharmacology. 2024 Aug;32(4):2125–2151. doi: 10.1007/s10787-024-01492-1.
29. Chandran B, Goel A. A randomized, pilot study to assess the efficacy and safety of curcumin in patients with active rheumatoid arthritis. Phytother Res. 2012;26(11):1422-1427. doi:10.1002/ptr.4639.
30. Arreola R, Quintero-Fabián S, López-Roa RI, Flores-Gutiérrez EO, Reyes-Grajeda JP, Carrera-Quintanar L, et al. Immunomodulation and anti-inflammatory effects of garlic compounds. J Immunol Res. 2015;2015:401630. doi:10.1155/2015/401630.
31. Qurat ul Ain. “The impact of climate change on preproduction and postproduction activities of food systems and vice versa.” Practices in Science and Technology (2025), 1, 1, 20-27. https://doi.org/10.56770/pst2025113
32. Mashhadi NS, Ghiasvand R, Askari G, Feizi A, Hariri M. Anti-oxidative and anti-inflammatory effects of ginger in health and physical activity: review of current evidence. Int J Prev Med. 2013;4(Suppl 1):S36-S42. doi:10.4103/2008-7802.116709.
33. Ryan JL, Heckler CE, Roscoe JA, Dakhil SR, Kirshner J, Flynn PJ, et al. Ginger (Zingiber officinale) reduces acute chemotherapy-induced nausea: a URCC CCOP study of 576 patients. Support Care Cancer. 2012;20(7):1479-1489. doi:10.1007/s00520-011-1236-3.
34. Possemiers S, Bolca S, Verstraete W, Heyerick A. The intestinal microbiome: a separate organ inside the body with the metabolic potential to influence the bioactivity of botanicals. Fitoterapia. 2011;82(1):53-66. doi:10.1016/j.fitote.2010.07.012.
35. Namba K, Hatano M, Yaeshima T, Takase M, Suzuki K. Effects of Bifidobacterium longum BB536 administration on influenza infection, influenza vaccine antibody titer, and cell-mediated immunity in the elderly. Biosci Biotechnol Biochem. 2010;74(5):939-945. doi:10.1271/bbb.90749.
36. Macpherson H, Pipingas A, Pase MP. Multivitamin-multimineral supplementation and mortality: a meta-analysis of randomized controlled trials. Am J Clin Nutr. 2013;97(2):437-444. doi:10.3945/ajcn.112.049304.
37. Chiva-Blanch G, Urpi-Sarda M, Ros E, Valderas-Martínez P, Casas R, Arranz S, et al. Effects of red wine polyphenols and alcohol on glucose metabolism and the lipid profile: a randomized clinical trial. Clin Nutr. 2013;32(2):200-206. doi:10.1016/j.clnu.2012.08.022.
38. Saad AM, Mohammed DM, Alkafaas SS, Ghosh S, Negm SH, Salem HM, Fahmy MA, Semary HE, Ibrahim EH, AbuQamar SF, El-Tarabily KA, et al. Dietary polyphenols and human health: sources, biological activities, nutritional and immunological aspects, and bioavailability—a comprehensive review. Front Immunol. 2025;16:1653378. doi: 10.3389/fimmu.2025.1653378